

1 **To:** Biofouling

2 *Mini Review*

3

4 **Title:** The oceans are changing: impact of ocean warming and acidification on biofouling
5 communities

6

7 **Authors:** Sergey Dobretsov^{1 2*}, Ricardo Coutinho³, Daniel Rittschof⁴, Maria Salta⁵,
8 Frederica Ragazzola⁵, Claire Hellio⁶

9

10 ¹Marine Science and Fisheries Department, College of Agricultural and Marine Sciences,
11 Sultan Qaboos University, Al-Khoud 34. PO Box 123, Sultanate of Oman.

12 ²Centre of Excellence in Marine Biotechnology, Sultan Qaboos University, Al-Khoud 50.
13 PO Box 123, Sultanate of Oman.

14 ³Instituto de Estudos do Mar Almirante Paulo Moreira, Rua Kioto nº 253, Praia dos Anjos,
15 Arraial do Cabo, RJ, Brazil.

16 ⁴Nicholas School, Duke University Marine Laboratory, 135 Duke Marine Lab Road,
17 Beaufort NC 28516 USA.

18 ⁵School of Biological Sciences, University of Portsmouth, King Henry Building,
19 Portsmouth, PO1 2DY, UK.

20 ⁶Université de Bretagne Occidentale, BIODIMAR/LEMAR UMR 6539, IUEM -
21 Technopole Brest-Iroise, rue Dumont d'Urville, 29280, Plouzané, France.

22 Total: 6,708 words

23 Without references: 3,672 words

24

Abstract

Climate change (CC) is driving modification in the chemical and physical properties of estuaries and oceans with profound consequences for species and ecosystems. Numerous studies investigate its effect from species to ecosystem levels, however little is known on impacts on biofilm communities and bioactive molecules, like cues, glues, and enzymes. CC is induced by anthropogenic activity increasing greenhouse emissions leading to rises in air and water temperatures, ocean acidification, sea level rise and changes in ocean gyres and rainfall patterns. These environmental changes are resulting in alterations in marine communities and spreading of species (pathogens, invasives). This review provides insights and synthesis of knowledge about the effect of elevated temperature and ocean acidification on microfouling communities and bioactive molecules. The existing studies suggest that CC will impact production of bioactive compounds, growth and composition of biofouling communities. Undoubtedly, with CC fouling management will become an even greater challenge.

Keywords: Biofilm, Biofouling, Bioactive compounds, Climate change

1. Introduction

The carbon dioxide concentration in the Earth's atmosphere is clearly and steadily rising (IPCC 2013). Anthropogenic emission has driven CO₂ concentration in the atmosphere from 280 ppm during the pre-industrial era to well over 400 ppm at the Hawaii monitoring site since 2015 with an estimated increase of 2ppm per year (<https://www.esrl.noaa.gov/gmd/ccgg/trends/weekly.html>). This increase in CO₂ concentration in the atmosphere causes many physical consequences for marine environments including ocean warming (IPCC 2013). The mean global sea surface temperature is increasing at an average rate of >0.1°C per decade (over the last 39 years), with the strongest warming trends found at high latitudes, and with future prediction estimating an increase of 2.7 °C by 2090. Temperature variations are often accompanied by changes in salinity due to reduced or enhanced precipitation relative to evaporation. Freshening and warming cause enhanced density stratification (IPCC 2013) and reduce the depth of winter mixing, which can cause a decrease in O₂ concentration in Oxygen Minimum Zones which are higher than the estimated decrease in O₂ concentration in the Open Ocean (mean rate in the of 0.1 to >0.3 μmol Kg⁻¹yr⁻¹) (IPCC 2013). Additionally, climate change is expected to increase upwelling frequency and intensity, lead to sea level rise due to melting of sea ice and glaciers (Doney et al. 2012). Increase in the level of atmospheric CO₂ will lead to ocean acidification (Doney et al. 2009). Because the oceanic and atmospheric gas concentrations tend towards equilibrium, ~30% of the atmospheric CO₂ has been taken up by the oceans, decreasing average pH by ~0.1 pH_T unit and ultimately changing water chemistry. The observed decrease in pH_T corresponds to a 26% increase in hydrogen ion concentration of seawater (Feely et al.

2009). By 2100, pH is expected to change by -0.13 (421ppm under RCP2.6), -0.22 (538ppm under RCP4.5), -0.28 (670ppm under RCP6.0) and -0.42 pH_T unit (936ppm under RCP8.5). Some progress has been made to understand the consequences of changes in pH, carbonate CO₃²⁻, and the saturation state of CaCO₃ for marine organisms and ecosystems (IPCC 2013; Wahl et al. 2015). These chemical and physical changes have direct implications for physiological processes such as photosynthesis, calcification, growth rates and internal pH regulation in a wide range of organisms (McCoy and Ragazzola 2014, Nannini et al. 2015, Evans et al. 2017, Fabricius et al. 2017, Okazaki et al. 2017) which will lead in a disruption of marine ecosystems and a reduction of biodiversity (Hoegh-Guldberg et al. 2007, Milazzo et al. 2014, Beaugrand et al. 2015).

All industrial installations in estuaries, bays, seas and oceans, such as vessels, platforms, buoys, quickly develop biofouling, a community composed of micro- and macro-fouling organisms (Clare et al. 1992). Micro-fouling usually presents as a dynamic microbial biofilm, which is composed of various species of bacteria, microalgae and protozoa incorporated in a muco-polysaccharide matrix (Dobretsov 2010; Malaeb et al. 2013; Salta et al. 2013). Macro-fouling communities are complex, with barnacles, bryozoa, mussels, polychaetes and macroalgae being the most common (Richmond & Seed 1991; Zardus et al. 2008). In some cases, micro-fouling organisms produce chemical cues that induce or inhibit settlement of macro-fouling species (Crisp 1984; Dobretsov et al. 2006; Hadfield 2011; Qian et al. 2007; Rittschof 2017) while in others there is a little direct relationship between macro- and micro-fouling.

Biofouling has a huge economic impact on maritime industries (Callow & Callow 2002; Trepos et al. 2014). Biofouling clogs aquaculture nets, water intakes, heat exchangers and

88 reduces ship hull performance (Okamura et al. 2010; Schultz et al. 2011; Sievers et al.
89 2014). Moreover, biofouling increases corrosion, shear stress and drag, eventually leading
90 to higher fuel consumption (Schultz et al. 2011) and increased production of CO₂ and
91 carbon particulates.

92 There are numerous reports of the effect of single environmental factors associated with
93 climate change (CO₂ level, elevated temperatures and acidification) on individual benthic
94 species (Bamber 1990; Parker et al. 2011; Lane et al. 2013; Calosi et al. 2013; Peck et al.
95 2015). Some of these benthic species, like the blue mussel *Mytilus edulis*, are potent
96 biofouling species. In contrast, the percent of biofouling publications dealing with climate
97 change is quite low but increasing every year (Figure 1). Several publications report effects
98 of factors associated with climate change on micro- and macro-fouling communities on
99 inert substrates (Kim & Micheli 2013; Gladis-Schmacka et al. 2014; Peck et al. 2015) and
100 living hosts (Nasrolahi et al. 2012; Stratil et al. 2013; Saderne & Wahl 2013; Saha et al.
101 2014).

102 A significant proportion of the biofouling-related climate change literature addresses
103 invasive species (Stachowicz et al. 2002; Hellmann et al. 2008; Canning-Clode et al. 2011).
104 Invasive species can be introduced by ship fouling and in ballast water (Davidson et al.
105 2008, 2009; Sorte et al. 2010; Keller et al. 2011). Most biofouling-related climate change
106 literature deals with species (organismal level) or populations of individual species (Figure
107 2). Fewer researchers investigated potential impact of factors associated with climate
108 change on multispecies communities. The lowest number of publications report effects of
109 factors associated with climate change on signaling molecules and the biochemistry of

organisms (Poloczanska & Butler 2009) (Figure 2). The impact of climate change on microbial communities and the bioactive molecules they generate is understudied.

This review focuses on the impacts of elevated temperature and ocean acidification, on biofouling communities. Particular focus is on the effect of factors associated with climate change on bioactive molecules of fouling organisms and growth and composition of microbial communities. Finally, we suggest areas for fruitful future investigation and the implication of climate change on the antifouling industry.

2. Climate change and bioactive molecules from fouling organisms

Biologically active molecules are produced by all phyla of marine organisms and play important roles in signaling, communication, allelopathy (Mayer et al. 2013) and organization of marine communities (Browne et al. 1998; Hay 2009; Rittschof 2017). Chemical cues from bacteria, diatoms and fungi induce or inhibit settlement of invertebrate larvae and algal spores (Wieczorek et al. 1996; Zardus et al. 2008; Dobretsov et al. 2013). When released in the marine environment, most of these biologically active molecules are bio-transformed or biodegraded by microbes (Uroz et al. 2005; Moree et al. 2012). There is a straightforward relationship between increase in temperature and the half-life of biologically active molecules (Singh et al. 2004). Similarly, there is a positive relationship between the concentration of heterotrophic bacteria and the half-life of signal molecules (Decho et al. 2010). Elevated water temperatures due to climate change will stimulate growth of microorganisms and enhance biodegradation of cues as well as enhance synthesis of antimicrobial compounds by marine fouling organisms (Table 1).

Marine bacteria coordinate virulence, competence, conjugation, production of antibiotics, motility, and biofilm formation by quorum sensing (QS) (Miller & Bassler 2001; Waters & Bassler 2005; Williams 2007). QS is based on the production, release and detection of chemical signal molecules called autoinducers. Increased concentrations of these signals due to high bacterial population density lead to an alteration in gene expression that regulates bacterial physiological activities (Decho et al. 2011). One of the most common and studied class of QS signal molecules is acyl homoserine lactone (AHL) (Waters & Bassler 2005). AHLs are unstable at >pH 7 (Yates et al. 2002). Studies assessing the stability of AHL against alkaline hydrolysis showed that AHLs having longer acyl chains (>12 carbons) are more resistant to breakdown than their shorter counterparts (Hmelo et al. 2011). In laboratory and field experiments, pH has a significant impact on the concentration of AHLs in microbial mats (Decho et al. 2009). In phototrophic microbial mats, short chain AHLs degrade quickly during the day, when the pH is > 8.2. During the night, when pH is 6.8 the concentrations of AHLs increases (Decho et al. 2009). When shorter-chain AHLs are degraded too rapidly, cellular communication may be disrupted. Acidification due to climate change will have a dramatic effect on concentrations of AHLs (Table 1). Since AHLs are important for biofilm structure and composition and settlement of some macro-fouling species (Dobretsov et al. 2009), it is possible that changes in production of QS compounds will alter densities and compositions of biofouling communities.

Enzymes are biological catalysts that accelerate the rate of specific biochemical reactions. Most enzymes are proteins and their structure is important for their activity. Increased temperature and changes in pH can lead to partial inhibition and in extreme cases to

inactivation of enzymes (Iyer & Ananthanarayan 2008). However, in other cases such as the activity of trypsin-like enzymes (Rittschof, 2017) increased temperature and lowered pH are near the optimum for the enzymes and increase rates of reactions. When marine organisms are subjected to environmental change (Hochachka & Somero 2002), the three main mechanisms used to maintain physiological homeostasis are: 1) quantitative (changing the concentration of enzymes and/or reactants); 2) qualitative (using a protein variant); 3) modulation (modifying the protein environment to reduce the impact of environmental change) (Clarke 2003).

Research shows that temperature impacts the enzyme levels and physiology of barnacles (Wong et al. 2011). Water temperature and high anthropogenic pollution have a significant effect on concentrations of antioxidant enzymes, such as catalase, superoxide dismutase and NADH-DT diaphorase, in the barnacle *Balanus* (= *Amphibalanus*) *amphitrite* (Niyogi et al. 2001). Anthropogenic ocean acidification alters protein expression patterns in *B. amphitrite* (Wong et al. 2011) although past studies have not found effects on reproduction due to changes in pH (McDonald et al. 2009, Nardone et al., 2018). The impact of temperature and pH on adhesion of barnacles has been reported (Tedesco et al. 2017).

Similarly, enzymes responsible for calcification of sedentary polychaete tubes were affected by elevated concentrations of CO₂ (Chan et al. 2012; Lane et al. 2013). Past work indicates the aragonite-producing juveniles of *Hydroides elegans* at the level of acidification predicted for the years 2050-2300 will not be able to maintain integrity of their calcification products (Chan et al. 2012).

Acidification affects interactions between iron and 3,4-dihydroxyphenylalanine (DOPA) and thus weakens byssus attachment of *Mytilus trossulus* to non-calcified materials

(O'Donnell et al. 2013). Mussel byssus threads were weaker and less extensible when secreted under elevated pCO₂ (>1200 µatm), whereas shell and tissue growth were unaffected (O'Donnell et al. 2013). Byssal fiber performance was reduced by 40%, which suggest that mussels will be dislodged by forces lower than those which dislodge them under present conditions. Decreased mussel attachment strength due to low pH was also reported by Zhao et al. (2017) who showed with real time PCR that low pH altered the expression of genes encoding proximal thread matrix protein, precursor collagen proteins and mussel foot proteins. The expression of some genes was down regulated, while others were up regulated. In multi-species communities, the impact of ocean acidification on mussel biomolecules became less predictable. A recent mesocosm study suggested that dense populations of macrophytes, like *Fucus vesiculosus* and *Zostera marina*, may mitigate acidification impact on mussel (*Mytilus edulis*) calcification by raising mean pH of seawater (Wahl et al. 2017). In the future, factors associated with climate change can change activity of enzymes and other bioactive molecules and, thus, change physiology and behavior of fouling organisms, and, finally, lead to changes in biofouling communities.

3. Climate change and microbial communities

Stress factors associated with climate change affect the growth and productivity of microbes (Rajkumar et al. 2013) and production of bioactive compounds (Hasegawa et al. 2005; Yang et al. 2007). Temperature has a dramatic impact on microbial growth (Price & Sowers 2004). Elevated temperature accelerates the growth of mesophiles and slows the growth of psychrophiles and alters the interactions between bacteria and their hosts (White et al. 1991; Wahl et al. 2012). In a case of marine pathogens, elevated temperature increases

growth, virulence and antimicrobial resistance (Kimes et al. 2012; Abdallah et al. 2014).
 For example, at 28⁰ C the infection rate and attachment of the coral pathogen *Vibrio shiloi*
 increases, while at the lower temperatures (about 16⁰ C) bacterial adhesion and growth in
 the tissues of the host coral *Oculina patagonica* is minimal and does not cause bleaching
 (Toren et al. 1998; Kushmaro et al. 2001). Virulence factors involved in motility, host
 degradation, secretion, antimicrobial resistance and transcriptional regulation are up-
 regulated in the pathogen *Vibrio coralliilyticus* at temperatures above 27⁰ C (Kimes et al.
 2012).
 Factors associated with climate change (e.g. increase in temperature, frequency of El-Nino
 and La-Nina-like conditions) and anthropogenically induced eutrophication cause massive
 algal blooms of microorganisms (Paerl & Huisman 2009). Due to presence of algal toxins
 and elevated oxygen consumption these blooms result in massive benthic and fish kills
 (Richlen et al. 2010; Hallegraeff 2010) and create estuarine and ocean dead zones (Diaz &
 Rosenberg 2008). In early July 2008, high level of nutrients and surface temperatures
 triggered a very dramatic bloom of *Ulva* sp. occurred in the China Sea off Qingdao, China
 (Leliaert et al. 2009). Similarly, in January-February 2014 extremely high ocean
 temperatures on the Atlantic coast of Brazil stimulated the largest algal bloom in the
 country's history. The bloom was composed of several species with the red alga
Aglaothamnion uruguayense being the most abundant (Martins et al. 2016). In 2008-2009
 in the Persian Gulf, an algal bloom of the dinoflagellate *Margalefidinium* (*Cochlodinium*)
polycricoides probably brought by ballast waters caused high mortality among benthic
 animals and fishes (Richlen et al. 2010) and dramatically decreased biomass of biofouling
 communities (Dobretsov 2015). These examples suggest that algal bloom conditions are

becoming the norm for most populated coastal regions and their impact on benthic and fouling community ecosystems will intensify in the warming oceans.

Marine biofilms are communities composed of viruses, bacteria, microalgae and protozoa incorporated in an exopolymer matrix (Zobell & Allen 1935; Webb et al. 2003; Qian et al. 2007; Dobretsov 2010). Biofilms are dynamic and the composition of communities can be altered by changes in environmental conditions, such as temperature, salinity, pH, and nutrient availability (Qian et al. 2007; Salta et al. 2013). For example, the number of rainy days and temperature affected growth of phototrophic biofilms on roof tiles (Gladis-Schmacka et al. 2014). Researchers studied the effect of different temperatures (high, low and ambient) on formation of microbial biofilms and subsequent larval settlement in laboratory experiments (Lau et al. 2005; Whalan & Webster 2014). Increased water temperatures led to formation of different microbial communities and subsequently affected settlement of larvae. Compositions of microbial communities associated with the alga *Fucus vesiculosus* were different when these algae were exposed to different temperatures or light intensities (Saha et al. 2014). Changes in pH led to significant decreases in biofilm performance and diversity (Patil et al. 2011). Peck and co-authors (Peck et al. 2015) studied formation of biofouling communities under ambient (pH = 7.9) and acidified (pH = 7.7) conditions at a constant temperature (23°C). After 100 days in acidified conditions, the proportion of sponges and ascidians is increased but numbers of the spirorbid *Neodexiospira pseudocorrugata* were reduced 5-fold. Changes in pH affected microfouling communities as well; the densities of the diatoms were lower in the low pH treatments compared to controls (Peck et al. 2015). Similarly, the microbial communities of corals, coralline algae and foraminifera were significantly different after the exposure to

pH 7.9 ($p\text{CO}_2 = 822 \mu\text{atm}$) over 6 weeks (Webster et al. 2013). In contrast, elevated $p\text{CO}_2$ had no impact on the microbiome associated with rhodoliths (Cavalcanti et al. 2018). If one ventures beyond the host algal thresholds to climate change, positive host-microbiome interactions are disrupted. Increasing temperatures resulted in a 2-fold increase in relative abundance of epibiotic Rhodobacteraceae on the surface of *F. vesiculosus* (Stratil et al. 2013). Similarly, community diversity measured by evenness and richness was higher at ambient water temperatures than at elevated temperatures. Thus, climate change can shift the structure of biofilms on inert and natural substrata (Table 1).

Biofilms play an important role by inducing or suppressing settlement of spores and larvae of some macrofouling species (Dobretsov et al. 2006; Zardus et al. 2008; Hadfield 2011; Salta et al. 2013). Thus, changes in microbial communities due to climate change could alter the structure of macro-fouling communities. For example, in the laboratory microbial communities developed at 23°C and 30°C were different from ones at 16°C (Lau et al. 2005). Larval response to these biofilms was also different; biofilms developed in the laboratory at 23°C and 30°C stimulated settlement of larvae of the barnacle *B. (=A.) amphitrite* and *B. trigonus* but had no effect on the polychaete larvae *Hydroides elegans* (Lau et al. 2005). Similarly, biofilms developed at elevated temperatures stimulated sponge larval settlement (Whalan & Webster 2014). Changes in the microbial community associated with crustose coralline algae reduced coral larval settlement under low pH (Webster et al. 2013). UV radiation reduces densities of bacteria in biofilms, which in turn decrease settlement of *Hydroides elegans* (Dobretsov et al. 2005). These examples show that temperature and pH associated with climate change directly affect composition and

densities of microorganisms in biofilms and indirectly (through biofilm composition and cues) reduce or enhance larval settlement of macro-fouling species.

4. Conclusions and future research directions

Climate change and increased anthropogenic activity will have strong effects on micro- and macro-fouling communities (Figure 3; Table 1). Though there are some publications on impacts of temperature, pH and generation of bioactive molecules, cues and signals associated with climate change at the species level, there is far less information about the impact of these factors at the community and molecular levels (Figure 2). This review suggested that increased temperatures and ocean acidification can affect compound production, detection, turnover and, in turn, will have a dramatic effect on microbial and macro-fouling communities.

Reports of the impact of ocean acidification on biofouling communities and their bioactive compounds are contradictory, indicating that responses are community dependent. Acidification will impact aragonite and magnesium calcite producers, such as coralline algae, corals, mussels, barnacles and some bryozoans (Doney et al. 2012; Chan et al. 2012; Lane et al. 2013). Acidified conditions significantly change biofouling community composition by a decrease in calcified (tube worms) and an increase in soft-bodied organisms, like ascidians and sponges (Peck et al. 2015). In contrast, some biofouling species (like *Amphibalanus amphitrite* and *Alcyonidium hirsutum*), their larvae and proteins are not sensitive to predicted changes in pH (McDonald et al. 2009; Saderne & Wahl 2013; Nardone et al., 2018). Moreover, dense populations of macroalgae, like *F. vesiculosus*, may reduce adverse effect of acidification on calcified biofouling organisms

(Wahl et al. 2017). Thus, it is likely that ecological impacts of ocean acidification will be location, species and community specific (Ekstrom & Moser 2014; Ekstrom et al. 2015). Future studies will answer questions about biofouling communities facing ocean acidification.

With climate change fouling management is a challenge (Table 2; Dobretsov 2009). Climate change will affect rates of leaching and dissolution of toxic ions and hydrolysis of copolymers of antifouling coatings because these are temperature, pH and flow dependent (Yebra et al. 2004; Yebra et al. 2006). Because coating chemistry and release rates are temperature sensitive, meeting environmental regulations in regions, which experience extreme temperatures, will be challenging. Additionally, spreading of invasive species (Sorte et al. 2010) will provide new challenges for industry. Novel regulations that will require coating companies to address these issues and provide new environmentally safe products that are effective in managing fouling in a warming and changing world are urgently needed.

As the polar ice melts, fast and inexpensive polar shipping routes are becoming possible (Lasserre & Pelletier 2011). In the future goods will travel on ships through the Arctic to Europe and Asia. However, the potential impacts of these new routes with respect to introduced and invasive species and performance of antifouling coatings remains unclear (Bax et al. 2003; Ware et al. 2014, Table 2). Long term information on biofouling communities in Arctic and preventive measures are lacking (Zvyagintsev 2003). Several important questions arise: Which invasive species have a chance to establish in warming Arctic waters? Will polar port biofouling communities develop that are comparable to those found in temperate and tropical regions? Will antifouling coatings designed for

temperate waters be effective and environmentally benign in warming Arctic? All of these questions should be answered urgently.

To conclude, we are at the beginning of our understanding of impacts of factors associated with climate change on bioactive molecules. The few existing studies suggest that ocean warming and acidification will have dramatic consequences on biofouling communities and their bioactive compounds. Probably, this effect will be region, community and species specific, which should be priority of future studies.

Acknowledgements

RC acknowledges CNPq Science without Frontier program for SD and CH, and CNPq for DR travel grant, CNPq for the Research Productivity Fellowship and INCT-PRO-OCEANO program. Part of the work of SD was supported by the HM Sultan Qaboos Research Trust Fund SR/AGR/FISH/10/01, the TRC grant RC/AGR/FISH/16/01 and a collaborative grant CL/SQU-SA/18/01. CH and RC acknowledge the EU LEAF (Low Emission AntiFouling) FP7 European project 314697 for financial support.

References

- Abdallah M, Chataigne G, Ferreira-Theret P, Benoliel C, Drider D, Dhulster P, Chihib N-E. 2014. Effect of growth temperature, surface type and incubation time on the resistance of *Staphylococcus aureus* biofilms to disinfectants. *Appl Microbiol Biotechnol*. 98:2597–2607.
- Bamber RN. 1990. The effects of acidic seawater on three species of lamellibranch mollusc. *J Exp Mar Biol Ecol*. 143:181–191.
- Bax N, Williamson A, Agüero M, Gonzalez E, Geeves W. 2003. Marine invasive alien species: a threat to global biodiversity. *Mar Policy*. 27:313–323.
- Beaugrand G, Edwards M, Raybaud V, Goberville E, Kirby RR. 2015. Future vulnerability of marine biodiversity compared with contemporary and past changes. *Nature Climate Change* 5:695.
- Browne KA, Tamburri MN, Zimmer-Faust RK. 1998. Modelling quantitative structure-activity relationships between animal behaviour and environmental signal molecules. *J Exp Biol*. 201:245–258.
- Callow ME, Callow JE. 2002. Marine biofouling: a sticky problem. *Biol Lond Engl*. 49:10–14.
- Calosi P, Rastrick SPS, Lombardi C, Guzman HJ de, Davidson L, Jahnke M, Giangrande A, Hardege JD, Schulze A, Spicer JJ, Gambi M-C. 2013. Adaptation and acclimatization to ocean acidification in marine ectotherms: an in situ transplant experiment with polychaetes at a shallow CO₂ vent system. *Philos Trans R Soc B Biol Sci*. 368:20120444.
- Canning-Clode J, Fowler AE, Byers JE, Carlton JT, Ruiz GM. 2011. “Caribbean Creep” Chills Out: Climate Change and Marine Invasive Species. *PLoS ONE*. 6:e29657.
- Cavalcanti GS, Shukla P, Morris M, Ribeiro B, Foley M, Doane MP, Thompson CC, Edwards MS, Dinsdale EA, Thompson FL. 2018. *Rhodoliths holobionts* in a changing ocean: host-microbes interactions mediate coralline algae resilience under ocean acidification. *BMC Genomics*. 19(1):701. doi: 10.1186/s12864-018-5064-4.
- Chan VBS, Li C, Lane AC, Wang Y, Lu X, Shih K, Zhang T, Thiyagarajan V. 2012. CO₂-Driven Ocean Acidification Alters and Weakens Integrity of the Calcareous Tubes Produced by the Serpulid Tubeworm, *Hydroides elegans*. *PLoS ONE*. 7:e42718.
- Clare AS, Rittschof D, Gerhart DJ, Maki JS. 1992. Molecular approaches to nontoxic antifouling. *Invertebr Reprod Dev*. 22:67–76.
- Clarke A. 2003. Costs and consequences of evolutionary temperature adaptation. *Trends Ecol Evol*. 18:573–581.

- 367 Crisp DJ. 1984. Overview of research on marine invertebrate larvae, 1940-1980. In:
368 Marine Biodeterioration: an Interdisciplinary Study. Editors J.D. Costlow and R.C. Tipper.
369 Naval Institute Press. Annapolis MD, pp 103-126.
370
- 371 Davidson IC, Brown CW, Sytsma MD, Ruiz GM. 2009. The role of containerships as
372 transfer mechanisms of marine biofouling species. *Biofouling*. 25:645–655.
- 373 Davidson IC, McCann LD, Sytsma MD, Ruiz GM. 2008. Interrupting a multi-species
374 bioinvasion vector: the efficacy of in-water cleaning for removing biofouling on obsolete
375 vessels. *Mar Pollut Bull*. 56:1538–1544.
- 376 Decho AW, Visscher PT, Ferry J, Kawaguchi T, He L, Przekop KM, Norman RS, Reid
377 RP. 2009. Autoinducers extracted from microbial mats reveal a surprising diversity of N-
378 acylhomoserine lactones (AHLs) and abundance changes that may relate to diel pH.
379 *Environ Microbiol*. 11:409–420.
- 380 Decho AW, Norman RS, Visscher PT. 2010. Quorum sensing in natural environments:
381 emerging views from microbial mats. *Trends Microbiol*. 18:73–80.
- 382 Decho AW, Frey RL, Ferry JL. 2011. Chemical challenges to bacterial AHL signaling in
383 the environment. *Chem Rev*. 111:86–99.
- 384 Diaz RJ, Rosenberg R. 2008. Spreading Dead Zones and Consequences for Marine
385 Ecosystems. *Science*. 321:926–929.
- 386 Dobretsov S. 2009. Expected effect if climate change on fouling communities and its
387 impact on antifouling research. In: Hellio C, Yebra D, editors. *Advances in marine*
388 *antifouling coatings and technologies*. Woodhead Publishing, CRC, Cambridge, UK, pp.
389 222-257
- 390 Dobretsov S. 2010. Marine Biofilms. In: Dürr S, Thomason JC, editors. *Biofouling*:
391 Wiley-Blackwell, p. 123–136.
- 392 Dobretsov S. 2015. Biofouling on artificial substrata in Muscat waters. *J Agr Mar Sci*. 20:
393 24-29.
- 394 Dobretsov SV, Qian P, Wahl M. 2005. Effect of solar ultraviolet radiation on the
395 formation of shallow, early successional biofouling communities in Hong Kong. *Mar*
396 *Ecol Prog Ser*. 290:55–65.
- 397 Dobretsov S, Dahms H-U, Qian P-Y. 2006. Inhibition of biofouling by marine
398 microorganisms and their metabolites. *Biofouling*. 22:43–54.
- 399 Dobretsov S, Teplitski M, Paul V. 2009. Mini-review: quorum sensing in the marine
400 environment and its relationship to biofouling. *Biofouling*. 25:413–427.
- 401 Dobretsov S, Abed RMM, Teplitski M. 2013. Mini-review: Inhibition of biofouling by
402 marine microorganisms. *Biofouling*. 29:423–441.

403 Doney SC, Fabry VJ, Feely RA, Kleypas JA. 2009. Ocean Acidification: The Other CO₂
404 Problem. *Annu Rev Mar Sci.* 1:169–192.

405 Doney SC, Ruckelshaus M, Emmett Duffy J, Barry JP, Chan F, English CA, Galindo
406 HM, Grebmeier JM, Hollowed AB, Knowlton N, et al. 2012. Climate Change Impacts on
407 Marine Ecosystems. *Annu Rev Mar Sci.* 4:11–37.

408 Ekstrom JA, Moser SC. 2014. Identifying and overcoming barriers in urban climate
409 adaptation: Case study findings from the San Francisco Bay Area, California, USA.
410 *Urban Clim.* 9:54–74.

411 Ekstrom JA, Suatoni L, Cooley SR, Pendleton LH, Waldbusser GG, Cinner JE, Ritter J,
412 Langdon C, van Hooijdonk R, Gledhill D, et al. 2015. Vulnerability and adaptation of US
413 shellfisheries to ocean acidification. *Nat Clim Change.* 5:207–214.

414 Evans TG, Pespeni MH, Hofmann GE, Palumbi SR, Sanford E. 2017. Transcriptomic
415 responses to seawater acidification among sea urchin populations inhabiting a natural pH
416 mosaic. *Molecular Ecology* 26:2257–2275.

417 Fabricius KE, Noonan SHC, Abrego D, Harrington L, De'ath G. 2017. Low recruitment
418 due to altered settlement substrata as primary constraint for coral communities under
419 ocean acidification. *Proceedings of the Royal Society B: Biological Sciences* 284.

420 Feely RA, Doney SC, Cooley SR. 2009. Ocean Acidification: Present Conditions and
421 Future Changes in a High-CO₂ world. *Oceanography.* 22:36–47.

422 Gladis-Schmack F, Glatzel S, Karsten U, Böttcher H, Schumann R. 2014. Influence of
423 local climate and climate change on aeroterrestrial phototrophic biofilms. *Biofouling.*
424 30:401–414.

425 Hadfield MG. 2011. Biofilms and Marine Invertebrate Larvae: What Bacteria Produce
426 That Larvae Use to Choose Settlement Sites. *Annu Rev Mar Sci.* 3:453–470.

427 Hallegraeff GM. 2010. Ocean Climate Change, Phytoplankton Community Responses,
428 and Harmful Algal Blooms: A Formidable Predictive Challenge1. *J Phycol.* 46:220–235.

429 Hasegawa H, Chatterjee A, Cui Y, Chatterjee AK. 2005. Elevated temperature enhances
430 virulence of *Erwinia carotovora* subsp. *carotovora* strain EC153 to plants and stimulates
431 production of the quorum sensing signal, N-acyl homoserine lactone, and extracellular
432 proteins. *Appl Environ Microbiol.* 71:4655–4663.

433 Hay M. 2009. Marine Chemical Ecology: Chemical signals and cues structure marine
434 populations, communities, and ecosystems. *Annual Review Mar Sci.* 1: 193–212.

435 Hellmann JJ, Byers JE, Bierwagen BG, Dukes JS. 2008. Cinco Consecuencias
436 Potenciales del Cambio Climático para Especies Invasoras. *Conserv Biol.* 22:534–543.

- 437 Hmelo LR, Mincer TJ, Van Mooy BAS. 2011. Possible influence of bacterial quorum
438 sensing on the hydrolysis of sinking particulate organic carbon in marine environments.
439 Environ Microbiol Rep. 3:682–688.
- 440 Hochachka PW, Somero GN. 2002. Biochemical adaptation: mechanism and process in
441 physiological evolution. Oxford University Press, 480p.
- 442 Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, Gomez E, Harvell
443 CD, Sale PF, Edwards AJ, Caldeira K, et al. 2007. Coral Reefs Under Rapid Climate
444 Change and Ocean Acidification. Science. 318:1737–1742.
- 445 IPCC 2013. Climate Change 2013: The Physical Science Basis. Contribution of Working
446 Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate
447 Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY,
448 USA.
- 449 Iyer PV, Ananthanarayan L. 2008. Enzyme stability and stabilization—Aqueous and non-
450 aqueous environment. Process Biochem. 43:1019–1032.
- 451 Keller RP, Drake JM, Drew MB, Lodge DM. 2011. Linking environmental conditions
452 and ship movements to estimate invasive species transport across the global shipping
453 network. Divers Distrib. 17:93–102.
- 454 Kimes NE, Grim CJ, Johnson WR, Hasan NA, Tall BD, Kothary MH, Kiss H, Munk AC,
455 Tapia R, Green L, et al. 2012. Temperature regulation of virulence factors in the
456 pathogen *Vibrio coralliilyticus*. ISME J. 6:835–846.
- 457 Kim TW, Micheli F. 2013. Decreased solar radiation and increased temperature combine
458 to facilitate fouling by marine non-indigenous species. Biofouling. 29:501–512.
- 459 Kushmaro A, Banin E, Loya Y, Stackebrandt E, Rosenberg E. 2001. *Vibrio shiloi* sp.
460 nov., the causative agent of bleaching of the coral *Oculina patagonica*. Int J Syst Evol
461 Microbiol. 51:1383–1388.
- 462 Lane AC, Mukherjee J, Chan VBS, Thiyagarajan V. 2013. Decreased pH does not alter
463 metamorphosis but compromises juvenile calcification of the tube worm *Hydroides*
464 *elegans*. Mar Biol. 160:1983–1993.
- 465 Lasserre F, Pelletier S. 2011. Polar super seaways? Maritime transport in the Arctic: an
466 analysis of shipowners' intentions. J Transp Geogr. 19:1465–1473.
- 467 Lau SCK, Thiyagarajan V, Cheung SCK, Qian P. 2005. Roles of bacterial community
468 composition in biofilms as a mediator for larval settlement of three marine invertebrates.
469 Aquat Microb Ecol. 38:41–51.
- 470 Leliaert F, Zhang X, Ye N, Malta E, Engelen AH, Mineur F, Verbruggen H, De Clerck
471 O. 2009. Research note: Identity of the Qingdao algal bloom. Phycol Res. 57:147–151.

- 472 Malaeb L, Le-Clech P, Vrouwenvelder JS, Ayoub GM, Saikaly PE. 2013. Do biological-
473 based strategies hold promise to biofouling control in MBRs? *Water Res.* 47:5447–5463.
- 474 Martins MS, Massocato TF, Horta PA, Barufi JB. 2016. First record of red macroalgae
475 bloom in Southern Atlantic Brazil. *Algae* 31(1): 33-39.
- 476 Mayer AMS, Rodríguez AD, Taglialatela-Scafati O, Fusetani N. 2013. Marine
477 pharmacology in 2009-2011: marine compounds with antibacterial, antidiabetic,
478 antifungal, anti-inflammatory, antiprotozoal, antituberculosis, and antiviral activities;
479 affecting the immune and nervous systems, and other miscellaneous mechanisms of
480 action. *Mar Drugs*. 11:2510–2573.
- 481 McCoy SJ, Ragazzola F. 2014. Skeletal trade-offs in coralline algae in response to ocean
482 acidification. *Nature Climate Change*. 4:719-723.
- 483 McDonald MR, McClintock JB, Amsler CD, Rittschof D, Angus RA, Orihuela B,
484 Lutostanski K. 2009. Effects of ocean acidification over the life history of the barnacle
485 *Amphibalanus amphitrite*. *Mar Ecol Prog Ser.* 385:179–187.
- 486 Milazzo M, Rodolfo-Metalpa R, Chan VBS, Fine M, Alessi C, Thiyagarajan V, Hall-
487 Spencer JM, Chemello R. 2014. Ocean acidification impairs vermetid reef recruitment.
488 *Scientific Reports* 4:4189.
- 489 Miller MB, Bassler BL. 2001. Quorum sensing in bacteria. *Annu Rev Microbiol.* 55:165–
490 199.
- 491 Moree WJ, Phelan VV, Wu C-H, Bandeira N, Cornett DS, Duggan BM, Dorrestein PC.
492 2012. Interkingdom metabolic transformations captured by microbial imaging mass
493 spectrometry. *Proc Natl Acad Sci.* 109:13811–13816.
- 494 Nannini M, De Marchi L, Lombardi C, Ragazzola F. 2015. Effects of thermal stress on
495 the growth of an intertidal population of *Ellisolandia elongata* (Rhodophyta) from N–W
496 Mediterranean Sea. *Mar Environ Res* 112, Part B:11-19.
- 497 Nardone JA, Patel S, Siegel KR, Tedesco D, McNicholl CG, O'Malley J, Herrick J,
498 Metzler RA, Orihuela B, Rittschof D, Dickinson GH. 2018. Assessing the Impacts of
499 Ocean Acidification on Adhesion and Shell Formation in the Barnacle *Amphibalanus*
500 *amphitrite*. *Frontiers in Marine Science*, section Marine Molecular Biology and Ecology.
501 <https://doi.org/10.3389/fmars.2018.00369>
- 502 Nasrolahi A, Stratil SB, Jacob KJ, Wahl M. 2012. A protective coat of microorganisms
503 on macroalgae: inhibitory effects of bacterial biofilms and epibiotic microbial
504 assemblages on barnacle attachment. *FEMS Microbiol Ecol.* 81:583–595.
- 505 Niyogi S, Biswas S, Sarker S, Datta AG. 2001. Seasonal variation of antioxidant and
506 biotransformation enzymes in barnacle, *Balanus balanoides*, and their relation with
507 polyaromatic hydrocarbons. *Mar Environ Res.* 52:13–26.

- 508 O'Donnell MJ, George MN, Carrington E. 2013. Mussel byssus attachment weakened by
509 ocean acidification. *Nat Clim Change*. 3:587–590.
- 510 Okamura D, Mori Y, Hashimoto T, Hori K. 2010. Effects of microbial degradation of
511 biofoulants on microfiltration membrane performance in a membrane bioreactor. *Environ*
512 *Sci Technol*. 44:8644–8648.
- 513 Okazaki RR, Towle EK, van Hooidonk R, Mor C, Winter RN, Piggot AM, Cuning R,
514 Baker AC, Klaus JS, Swart PK, Langdon C. 2017. Species-specific responses to climate
515 change and community composition determine future calcification rates of Florida Keys
516 reefs. *Global Change Biology* 23:1023-1035.
- 517 Paerl HW, Huisman J. 2009. Climate change: a catalyst for global expansion of harmful
518 cyanobacterial blooms. *Environ Microbiol Rep*. 1:27–37.
- 519 Parker LM, Ross PM, O'Connor WA. 2011. Populations of the Sydney rock oyster,
520 *Saccostrea glomerata*, vary in response to ocean acidification. *Mar Biol*. 158:689–697.
- 521 Patil SA, Harnisch F, Koch C, Hübschmann T, Fetzer I, Carmona-Martínez AA, Müller
522 S, Schröder U. 2011. Electroactive mixed culture derived biofilms in microbial
523 bioelectrochemical systems: The role of pH on biofilm formation, performance and
524 composition. *Bioresour Technol*. 102:9683–9690.
- 525 Peck LS, Clark MS, Power D, Reis J, Batista FM, Harper EM. 2015. Acidification effects
526 on biofouling communities: winners and losers. *Glob Change Biol*. 21(5):1907-1913.
- 527 Poloczanska ES, Butler AJ. 2009. Biofouling and Climate Change. In: Dürr S.,
528 Thomason J.C., editors. *Biofouling*, Wiley-Blackwell, p. 333–347.
- 529 Price PB, Sowers T. 2004. Temperature dependence of metabolic rates for microbial
530 growth, maintenance, and survival. *Proc Natl Acad Sci U S A*. 101:4631–4636.
- 531 Qian P-Y, Lau SCK, Dahms H-U, Dobretsov S, Harder T. 2007. Marine biofilms as
532 mediators of colonization by marine macroorganisms: implications for antifouling and
533 aquaculture. *Mar Biotechnol N Y N*. 9:399–410.
- 534 Rajkumar M, Prasad MNV, Swaminathan S, Freitas H. 2013. Climate change driven
535 plant-metal-microbe interactions. *Environ Int*. 53:74–86.
- 536 Richlen ML, Morton SL, Jamali EA, Rajan A, Anderson DM. 2010. The catastrophic
537 2008–2009 red tide in the Arabian gulf region, with observations on the identification and
538 phylogeny of the fish-killing dinoflagellate *Cochlodinium polykrikoides*. *Harmful Algae*.
539 9:163–172.
- 540 Richmond MD, Seed R. 1991. A review of marine macrofouling communities with
541 special reference to animal fouling. *Biofouling*. 3:151–168.

542 Rittschof D. 2017. Trypsins: Keystone Enzymes in Marine Communities. *Enzymology*
543 *and Protein Science. JSM Enzymol Protein Sci* 2(1): 1009

544 Saderne V, Wahl M. 2013. Differential Responses of Calcifying and Non-Calcifying
545 Epibionts of a Brown Macroalga to Present-Day and Future Upwelling pCO₂. *PLoS*
546 *ONE* 8(7): e70455. <https://doi.org/10.1371/journal.pone.0070455>

547 Saha M, Rempt M, Stratil SB, Wahl M, Pohnert G, Weinberger F. 2014. Defence
548 Chemistry Modulation by Light and Temperature Shifts and the Resulting Effects on
549 Associated Epibacteria of *Fucus vesiculosus*. *PLoS ONE*. 9:e105333.

550 Salta M, Wharton JA, Blache Y, Stokes KR, Briand J-F. 2013. Marine biofilms on
551 artificial surfaces: structure and dynamics. *Environ Microbiol*. 15:2879–2893.

552 Schultz MP, Bendick JA, Holm ER, Hertel WM. 2011. Economic impact of biofouling
553 on a naval surface ship. *Biofouling*. 27:87–98.

554 Sievers M, Dempster T, Fitridge I, Keough MJ. 2014. Monitoring biofouling
555 communities could reduce impacts to mussel aquaculture by allowing synchronisation of
556 husbandry techniques with peaks in settlement. *Biofouling*. 30:203–212.

557 Singh S, Singh A, Aziz MA, Waheed SM, Bhat R, Bhatnagar R. 2004. Thermal
558 inactivation of protective antigen of *Bacillus anthracis* and its prevention by polyol
559 osmolytes. *Biochem Biophys Res Commun*. 322:1029–1037.

560 Sorte CJB, Williams SL, Zerebecki RA. 2010. Ocean warming increases threat of
561 invasive species in a marine fouling community. *Ecology*. 91:2198–2204.

562 Stachowicz JJ, Terwin JR, Whitlatch RB, Osman RW. 2002. Linking climate change and
563 biological invasions: Ocean warming facilitates nonindigenous species invasions. *Proc*
564 *Natl Acad Sci*. 99:15497–15500.

565 Tedesco D, Nardone JA, Siegel KR, Patel S, Orihuela B, Rittschof D, Dickinson GH.
566 2017. Effects of ocean acidification and temperature on biomineralization and adhesion
567 in the barnacle, *Amphibalanus amphitrite*. Presented at 46th Benthic Ecology Meeting,
568 Myrtle Beach, South Carolina.

569 Trepos R, Pinori E, Berglin M, Svenson J, Coutinho R, Lausmaa J, Hellio C. 2014.
570 Innovative approaches for the development of new copper-free marine antifouling paints.
571 *Journal of ocean technology, J Ocean Technol*. 9:7–18

572 Toren A, Landau L, Kushmaro A, Loya Y, Rosenberg E. 1998. Effect of Temperature on
573 Adhesion of *Vibrio* Strain AK-1 to *Oculina patagonica* and on Coral Bleaching. *Appl*
574 *Environ Microbiol*. 64:1379–1384.

575 Uroz S, Chhabra SR, Cámara M, Williams P, Oger P, Dessaux Y. 2005. N-
576 Acylhomoserine lactone quorum-sensing molecules are modified and degraded by

577 *Rhodococcus erythropolis* W2 by both amidolytic and novel oxidoreductase activities.
578 Microbiol Read Engl. 151:3313–3322.

579 Wahl M, Goecke F, Labes A, Dobretsov S, Weinberger F. 2012. The second skin:
580 ecological role of epibiotic biofilms on marine organisms. Front Microbiol. 3:292.

581 Wahl M., Saderne V, Sawall Y. 2015. How good are we at assessing the impact of ocean
582 acidification in coastal systems? Limitations, omissions and strengths of commonly used
583 experimental approaches with special emphasis on the neglected role of fluctuations. Mar
584 Freshwater Res 67(1): 25-36.

585 Wahl M, Scheider Covacha S, Saderne V, Hiebenthal C, Muller JD, Pansch C, Sawall Y.
586 2017. Macroalgae may mitigate ocean acidification effects on mussel calcification by
587 increasing pH and its fluctuations. Limnology and Oceanography 63 (1): 3-21.

588 Ware C, Berge J, Sundet JH, Kirkpatrick JB, Coutts ADM, Jelmert A, Olsen SM, Floerl
589 O, Wisz MS, Alsos IG. 2014. Climate change, non-indigenous species and shipping:
590 assessing the risk of species introduction to a high-Arctic archipelago. Divers Distrib.
591 20:10–19.

592 Waters CM, Bassler BL. 2005. Quorum sensing: Cell-to-Cell Communication in
593 Bacteria. Annu Rev Cell Dev Biol. 21:319–346.

594 Webb JS, Givskov M, Kjelleberg S. 2003. Bacterial biofilms: prokaryotic adventures in
595 multicellularity. Curr Opin Microbiol. 6:578–585.

596 Webster NS, Negri AP, Flores F, Humphrey C, Soo R, Botté ES, et al. 2013. Near-Future
597 Ocean acidification causes differences in microbial associations within diverse coral reef
598 taxa. Environ Microbiol Rep. 5:243–251. doi: 10.1111/1758-2229.12006.

599 Whalan S, Webster NS. 2014. Sponge larval settlement cues: the role of microbial
600 biofilms in a warming ocean. Sci Rep 4.

601 White PA, Kalff J, Rasmussen JB, Gasol JM. 1991. The effect of temperature and algal
602 biomass on bacterial production and specific growth rate in freshwater and marine
603 habitats. Microb Ecol. 21:99–118.

604 Wieczorek SK, Murray AW, Todd CD. 1996. Seasonal variation in the effects of hard
605 substratum biofilming on settlement of marine invertebrate larvae. Biofouling. 10:309–
606 330.

607 Williams P. 2007. Quorum sensing, communication and cross-kingdom signalling in the
608 bacterial world. Microbiology. 153:3923–3938.

609 Wong KKW, Lane AC, Leung PTY, Thiagarajan V. 2011. Response of larval barnacle
610 proteome to CO₂-driven seawater acidification. Comp Biochem Physiol Part D
611 Genomics Proteomics. 6:310–321.

- 612 Yang LH, Miao L, Lee OO, Li X, Xiong H, Pang K-L, Vrijmoed L, Qian P-Y. 2007.
613 Effect of culture conditions on antifouling compound production of a sponge-associated
614 fungus. *Appl Microbiol Biotechnol.* 74:1221–1231.
- 615 Yates EA, Philipp B, Buckley C, Atkinson S, Chhabra SR, Sockett RE, Goldner M,
616 Dessaux Y, Cámara M, Smith H, Williams P. 2002. N-acylhomoserine lactones undergo
617 lactonolysis in a pH-, temperature-, and acyl chain length-dependent manner during
618 growth of *Yersinia pseudotuberculosis* and *Pseudomonas aeruginosa*. *Infect Immun.*
619 70:5635–5646.
- 620 Yebra DM, Kiil S, Dam-Johansen K. 2004. Antifouling technology—past, present and
621 future steps towards efficient and environmentally friendly antifouling coatings. *Prog Org*
622 *Coat.* 50:75–104.
- 623 Yebra DM, Kiil S, Weinell CE, Dam-Johansen K. 2006. Dissolution rate measurements
624 of sea water soluble pigments for antifouling paints: ZnO. *Prog Org Coat.* 56:327–337.
- 625 Zardus JD, Nedved BT, Huang Y, Tran C, Hadfield MG. 2008. Microbial biofilms
626 facilitate adhesion in biofouling invertebrates. *Biol Bull.* 214:91–98.
- 627 Zhao X, Guo C, Han Y, Che Z, Wang Y, Wang X. et al. 2017. Ocean acidification
628 decreases mussel byssal attachment strength and induces molecular byssal responses.
629 *Mar. Ecol. Prog. Ser.* 565: 67–77. doi: 10.3354/meps11992
- 630 Zobell CE, Allen EC. 1935. The Significance of Marine Bacteria in the Fouling of
631 Submerged Surfaces. *J Bacteriol.* 29:239–251.
- 632 Zvyagintsev AY. 2003. Introduction of Species into the Northwestern Sea of Japan and
633 the Problem of Marine Fouling. *Rus J Mar Biol* 29: 10-21.
- 634